

preventing the development of resistance in pest insects.

REFERENCES

- Bloomquist JR, Ion channels as target for insecticides. *Ann Rev Entomol* 41:163–190 (1996).
- Gordon D, Merrick D, Wollner DA and Catterall WA, Biochemical properties of sodium channels in a wide range of excitable tissues studied with site-directed antibodies. *Biochemistry* 27:7032–7038 (1988).
- Gordon D, Moskowitz H and Zlotkin E, Sodium channel polypeptides in central nervous systems of various insects identified with site-directed antibodies. *Biochim Biophys Acta* 1026:80–86 (1990).
- Loughney K, Kreber R and Ganetzky B, Molecular analysis of the *para* locus, a sodium channel gene in *Drosophila*. *Cell* 58:1143–1154 (1989).
- Moskowitz H, Herrmann R, Zlotkin E and Gordon D, Variability among insect sodium channels revealed by binding of selective neurotoxins. *Insect Biochem Molec Biol* 24:13–19 (1994).
- Gordon D, Martin-Eauclaire MF, Cestèle S, Kopeyan C, Carlier E, Ben Khalifa R, Pelhate M and Rochat H, Scorpion toxins affecting sodium current inactivation bind to distinct homologous receptor sites on rat brain and insect sodium channels. *J Biol Chem* 271:8034–8045 (1996).
- Fishman L, Herrmann R, Gordon D and Zlotkin E, Insect tolerance to neurotoxic polypeptide: pharmacokinetic and pharmacodynamic aspects. *J Exp Biol* 200:1115–1123 (1997).
- Zlotkin E, Miranda F and Rochat H, Chemistry and pharmacology of Buthinae scorpion venoms, in *Arthropod Venoms*, ed. by Bettini S, Springer Verlag pp 317–369 (1978).
- Lester D, Lazarovici P, Pelhate M and Zlotkin E, Purification, characterization and action of two insect toxins from the venom of the scorpion *Buthotus judaicus*. *Biochim Biophys Acta* 701:370–381 (1982).
- Eitan M, Fowler E, Herrmann R, Duval A, Pelhate M and Zlotkin E, A scorpion venom neurotoxin paralytic to insects that affects sodium current inactivation: purification, primary structure, and mode of action. *Biochemistry* 29:5941–5947 (1990).
- Zlotkin E, Kadouri D, Gordon D, Pelhate M, Martin MF and Rochat H, An excitatory and a depressant insect selective toxin from scorpion venom both affect sodium conductance and possess a common binding site. *Arch Biochem Biophys* 240:877–887 (1985).
- Gordon D, Jover E, Couraud F and Zlotkin E, The binding of an insect selective neurotoxin (AaIT) from scorpion venom to locust synaptosomal membranes. *Biochim Biophys Acta* 778:349–358 (1984).
- Gordon D, Moskowitz H, Eitan M, Warner C, Catterall WA and Zlotkin E, Localization of receptor sites for insect-selective toxins on sodium channels by site-directed antibodies. *Biochemistry* 31:7622–7628 (1992).
- Gordon D and Zlotkin E, Binding of an alpha scorpion toxin to insect sodium channels is not dependent on membrane potential. *FEBS Lett* 315:125–129 (1993).
- Zlotkin E, Eitan M, Bindokas V, Adams ME, Moyer M, Brukhart W and Fowler E, Functional duality and structural uniqueness of depressant insect-selective neurotoxins. *Biochemistry* 30:4814–4820 (1991).
- Gordon D, Savarin P, Gurevitz M and Zinn-Justin S, Functional anatomy of scorpion toxins affecting sodium channels. *J Toxicol Toxin Rev* 17:131–158 (1998).
- Zilberberg N, Gordon D, Pelhate M, Adams ME, Norris F, Zlotkin E and Gurevitz M, Functional expression and genetic modification of an alpha scorpion neurotoxin. *Biochemistry* 35:10215–10222 (1996).
- Zilberberg N, Froy O, Loret E, Cestèle S, Arad D, Gordon D and Gurevitz M, Identification of structural elements of a scorpion α -neurotoxin important for receptor site recognition. *J Biol Chem* 272:14810–14816 (1997).
- Tugarinov V, Kustanovich I, Zilberberg N, Gurevitz M and Anglister J, Solution structure of a highly insecticidal recombinant scorpion α -toxin and a mutant with increased activity. *Biochemistry* 36:2414–2424 (1997).
- Gordon D, Sodium channels as targets for neurotoxins: mode of action and interaction of neurotoxins with receptor sites on sodium channels, in *Toxins And Signal Transduction*, ed by Lazarowici P and Gutman Y, Harwood Press, Amsterdam. pp 119–149 (1997).
- Cestèle S, Ben Khalifa R, Pelhate M, Rochat H and Gordon D, α -Scorpion toxins binding on rat brain and insect sodium channels reveal divergent allosteric modulations by brevetoxin and veratridine. *J Biol Chem* 270:15153–15161 (1995).
- Oren DA, Froy O, Amit E, Kleinberger-Doron N, Gurevitz M and Shaanan B, An excitatory scorpion toxin with a distinctive feature: an additional α helix at the C terminus and its implications for interaction with insect sodium channels. *Structure* 6:1095–1103 (1998).
- Gordon D, A new approach to insect-pest control – combination of neurotoxins interacting with voltage-sensitive sodium channels to increase selectivity and specificity. *Invertebr Neurosci* 3:103–116 (1997).

Characterisation of the relationship between binding sites for imidacloprid and other nicotinic ligands in insects

Robert J Lind,^{1*} Martin S Clough,² Fergus GP Earley,² Susan Wonnacott¹ and Stuart E Reynolds¹

¹Department of Biology and Biochemistry, University of Bath, Claverton Down, Bath, BA2 7AY, UK

²Zeneca Agrochemicals, Jealott's Hill Research Station, Bracknell, Berkshire, RG42 6ET, UK

Abstract: Radioligand binding studies using the neurotoxins α -bungarotoxin, epibatidine, imidacloprid (IMI) and methyllycaonitine reveal heterogeneity at the level of the nicotinic acetylcholine receptor (nAChR) in membranes from the peach potato aphid *Myzus persicae* (Sulzer) and further suggest the presence of more than one ligand binding site per nAChR. These sites are able to interact allosterically with each other. Of particular interest, [³H]IMI has over an order of magnitude higher affinity in membranes of hemipteran pest species than in non-hemipteran insects, which may help explain why IMI is particularly effective for the control of sucking pests.

Keywords: *Myzus persicae*; nicotinic acetylcholine receptor; imidacloprid; methyllycaonitine; epibatidine; radioligand binding

* Correspondence to: Robert J Lind, Zeneca Agrochemicals, Jealott's Hill Research Station, Bracknell, Berks RG42 6ET, UK
E-mail: rob.lind@aguk.zeneca.com

Contract/grant sponsor: Biotechnology and Biological Science Research Council

Contract/grant sponsor: Zeneca Agrochemicals
(Received 10 August 1998; accepted 20 May 1999)

Table 1. Comparison of [3 H]IMI saturable binding in a panel of insects. Data for *Myzus persicae* is the mean (\pm SEM) of $n=5$, and $n=1$ for other insects

Insect ^a	High affinity		Low affinity		Hill value (n_H)
	K_d (nM)	B_{max} (fmolmg ⁻¹)	K_d (nM)	B_{max} (fmolmg ⁻¹)	
<i>M persicae</i>	0.14 (\pm 0.01)	284 (\pm 16)	12.58 (\pm 1.83)	883 (\pm 142)	0.61 (\pm 0.02)
<i>B tabaci</i>	0.10	19	2.1	112	0.50
<i>N cincticeps</i>	<0.01	33	1.23	179	0.67
<i>H virescens</i> *	1.51	134			0.92
<i>D melanogaster</i>	1.42	126			1.04
<i>P americana</i> *	3.14	2140			1.01
<i>C felis</i>	4.82	369			0.95

* Heads only.

1 INTRODUCTION

The naturally occurring neurotoxins α -bungarotoxin (α -BgTx), epibatidine (EPI) and methyllycaconitine (MLA), all known to interact specifically with vertebrate nicotinic acetylcholine receptors (nAChRs), were used to investigate nAChR of insects, principally the aphid *Myzus persicae* (Sulzer). The synthetic neonicotinoid insecticide imidacloprid (IMI), which is particularly effective against sucking insect pests such as aphids, was also studied.¹

2 EXPERIMENTAL METHODS

[3 H]EPI (55Ci mmol⁻¹), [3 H]IMI (30Ci mmol⁻¹), unlabelled IMI and [3 H] α -BgTx (60Ci mmol⁻¹) were supplied by Zeneca Agrochemicals. Na¹²⁵I, obtained from Amersham International (UK), was used to iodinate α -BgTx to a specific activity of 750Ci mmol⁻¹. [3 H]MLA (\sim 50 Ci mmol⁻¹) was synthesised by Tocris Cookson Ltd, Bristol, England. Other unlabelled drugs were obtained from Sigma-Aldrich Ltd (Poole, Dorset, UK).

Insect membranes were prepared as described elsewhere for whole peach potato aphid *M persicae*, cotton whitefly *Bemisia tabaci* (Gennadius), green leaf hopper *Nephotettix cincticeps* (Uhler), cat flea *Ctenocephalides felis* (Bouche), and fruitfly *Drosophila melanogaster* (Meigen), and heads of the tobacco budworm *Heliothis virescens* (F) and the cockroach *Periplaneta americana* (L).² Saturation, displacement and dissociation experiments employed either filtration or centrifugation to separate bound from free ligand using a cell harvester. Radioactivity was determined with scintillation spectrometry.² Isotopic dissociation studies employed either 1 μ M α -BgTx, EPI, IMI or MLA to initiate dissociation from *M persicae* membranes after an association period of 3 h with [3 H] α -BgTx. MLA was used at 1 μ M to determine non-specific binding for all experiments.

3 RESULTS

Saturable binding in *M persicae* membranes demonstrated that [125 I] α -BgTx ($K_d=1.2$ & 34 nM, $B_{max}=$




Figure 1. Isotopic dissociation of [3 H] α -BgTx in *M persicae* membranes initiated by either 1 μ M α -BgTx or MLA. Data shown are the mean of three experiments (\pm SEM).

167 & 640 fmolmg⁻¹), [3 H]EPI ($K_d=0.9$ & 18 nM, $B_{max}=344$ & 904 fmolmg⁻¹) and [3 H]IMI ($K_d=0.14$ & 13 nM, $B_{max}=284$ & 883 fmolmg⁻¹) each labelled two sites. In contrast, [3 H]MLA binding was resolvable as only a single binding site with a K_d of 0.95 nM and a B_{max} of 1290 fmolmg⁻¹. Similar studies using membranes of two hemipteran insects, *B tabaci* and *N cincticeps*, also revealed the presence of high- and low-affinity [3 H]IMI binding components. By contrast, studies in non-hemipteran insects could only resolve a single class of [3 H]IMI binding site. Of particular interest, the high affinity site labelled by [3 H]IMI in all of these hemipteran insects was at least an order of magnitude higher in affinity than was the single site in non-hemipteran insects (Table 1).

The rate of [3 H] α -BgTx isotopic dissociation was slow and monophasic from its high affinity binding site in *M persicae* membranes when initiated by either 1 μ M

α -BgTx, EPI or IMI. In contrast the dissociation rate was markedly increased when initiated by MLA compared to α -BgTx, EPI or IMI (Fig 1).

Displacement of [125 I] α -BgTx, [3 H]EPI and [3 H]IMI by MLA was with high potency with an IC₅₀ value close to the K_d value of each of these labelled ligands. Displacement of [3 H]MLA by α -BgTx, EPI and IMI resulted in low potency, but was characterised by shallow displacement curves.

4 DISCUSSION AND CONCLUSIONS

Saturation studies in *M persicae* membranes demonstrate that [125 I] α -BgTx, [3 H]EPI, and [3 H]IMI all label two binding components of differing affinities in a ratio of 1:3. In each case the sum of the high and low affinity B_{max} values approximated the B_{max} value of the single site labelled by [3 H]MLA. [3 H]MLA is a novel radioligand which is highly selective in vertebrate nAChRs,³ but which appears to be unable to distinguish between a heterogeneous population of nicotinic ligand binding sites in insects.

The increase in dissociation rate of [3 H] α -BgTx using MLA instead of either α -BgTx, EPI or IMI is consistent with an allosteric interaction by MLA at a binding site other than that of the high affinity [3 H] α -BgTx binding site.

Displacement studies have shown that MLA is a potent ligand at sites labelled with high affinity by [125 I] α -BgTx, [3 H]EPI, and [3 H]IMI. The shallow displacement curves obtained when [3 H]MLA is displaced by α -BgTx, EPI, and IMI (data not shown) are likely to be due to displacement of [3 H]MLA from at least two distinct binding sites, for which the displacing ligands have different affinities, but which are indistinguishable by [3 H]MLA.

Interestingly, the high affinity binding of [3 H]IMI in hemipteran membranes compared to that in non-hemipteran insects may help explain why IMI is particularly useful for the control of sucking pests.

ACKNOWLEDGEMENTS

Thanks to Matt Cahill of IACR Rothamsted for supplying *B tabaci*. The work was supported by the BBSRC and Zeneca Agrochemicals.

REFERENCES

- 1 Elbert A, Becker B, Hartwig J and Erdelen C, Imidacloprid – a new systemic insecticide. *Pflanzenschutz-Nachrichten Bayer* 44:113–136 (1991).
- 2 Lind RJ, Clough MS, Reynolds SE and Earley FGP, [3 H]Imidacloprid labels high and low affinity nicotinic acetylcholine receptor-like binding sites in the aphid *Myzus persicae* (Hemiptera: Aphididae). *Pestic Biochem Physiol* 62:3–14 (1998).
- 3 Davies ARL, Hardick DJ, Blagbrough IS, Potter BVL, Wolstenholme AJ and Wonnacott S, [3 H]Methyllycaconitine a novel radioligand for $\alpha 7$ -type nicotinic acetylcholine receptors. *Soc Neurosci Abstr* 23:477.2 (1997).

Nicotinic acetylcholine receptor chimeras of rat $\alpha 7$ and *Drosophila* SAD reveal species-specific agonist binding regions

Ingeborg van den Beukel,^{1*} Remco Klaassen,² Guus B Smit,² Regina GDM van Kleef¹ and Marga Oortgiesen^{1†}

¹Research Institute of Toxicology, Utrecht University, PO Box 80176, 3508 TD Utrecht, The Netherlands

²Department of Molecular Neurobiology, Graduate School of Neuroscience, Free University, De Boelelaan 1087, 1081 HV Amsterdam, The Netherlands

Abstract: Species-specific agonist binding regions of nicotinic acetylcholine receptors (nAChR) were examined. Imidacloprid and physostigmine (Phy) selectively activated insect nAChR composed of *Drosophila* second alpha-like subunit (SAD) and chick $\beta 2$, in contrast to rat $\alpha 7$ nAChR. The Phy-activated currents were α -bungarotoxin (α -BGT) sensitive, suggesting activation at the agonist binding loop C. Several SAD- $\alpha 7$ chimeras were constructed, by switching agonist binding regions, and expressed in oocytes. Though none of the chimeras was activated by a range of nicotinic agonists, [125 I] α -BGT binding revealed homomeric assembly of all chimeric cDNAs. Phy differentially displaced [125 I] α -BGT from the nAChR chimeras, suggesting that the β subunit is not involved in Phy binding, and that Phy targets the insect agonist binding loop C.

Keywords: nicotinic receptor; agonist binding; chimera; physostigmine; imidacloprid; acetylcholine receptor; neonicotinoid

1 INTRODUCTION

Multiple nicotinic acetylcholine receptor (nAChR) subtypes exist, both within and between species. Insects express only neuronal nAChR, while, in vertebrates, endplate and neuronal nAChR types are distinguished. A range of different insect and vertebrate α subunits exist, which may be combined with various β subunits to form functional nAChR. Three loops in the α subunits are thought to be involved in acetylcholine (ACh) binding,¹ ie loop A (residues 86–93), loop B (residues 148–151), and loop C (residues 190–198). The latter includes the two adjacent cysteines characteristic of α subunits. Distinct nAChR subtypes are distinguished by their physiological and pharmacological properties. In addition, differential sensitivities to chemical compounds may occur, suggesting a role in species-selective toxicity. The nitroguanidine insecticide imidacloprid selectively activates insect nAChR by binding to the agonist

* Correspondence to: Ingeborg van den Beukel, Research Institute of Toxicology, Utrecht University, PO Box 80176, 3508 Utrecht, The Netherlands
E-mail: I.vandenBeukel@ritox.dgk.ruu.nl

† Present address: Department of Neurobiology, Duke University Medical Center, 435 Bryan Research Building, Durham, NC 27278, USA

(Received 20 August 1998; revised version received 25 November 1998; accepted 20 May 1999)